
Parkin, PINK1, and DJ-1 form a ubiquitin E3 ligase complex promoting unfolded protein degradation.

Journal: J Clin Invest

Publication Year: 2009

Authors: Hui Xiong, Danling Wang, Linan Chen, Yeun Su Choo, Hong Ma, Chengyuan Tang, Kun Xia, Wei Jiang, Ze'ev Ronai, Xiaoxi Zhuang, Zhuohua Zhang

PubMed link: 19229105

Funding Grants: Modeling Parkinson's Disease Using Human Embryonic Stem Cells, Derivation of Parkinson's Disease Coded-Stem Cells (PD-SCs)

Public Summary:

Scientific Abstract:

Mutations in PARKIN, pten-induced putative kinase 1 (PINK1), and DJ-1 are individually linked to autosomal recessive early-onset familial forms of Parkinson disease (PD). Although mutations in these genes lead to the same disease state, the functional relationships between them and how their respective disease-associated mutations cause PD are largely unknown. Here, we show that Parkin, PINK1, and DJ-1 formed a complex (termed PPD complex) to promote ubiquitination and degradation of Parkin substrates, including Parkin itself and Synphilin-1 in neuroblastoma cells and human brain lysates. Genetic ablation of either Pink1 or Dj-1 resulted in reduced ubiquitination of endogenous Parkin as well as decreased degradation and increased accumulation of aberrantly expressed Parkin substrates. Expression of PINK1 enhanced Parkin-mediated degradation of heat shock-induced misfolded protein. In contrast, PD-pathogenic Parkin and PINK1 mutations showed reduced ability to promote degradation of Parkin substrates. This study identified a functional ubiquitin E3 ligase complex consisting of PD-associated Parkin, PINK1, and DJ-1 to promote degradation of un-/misfolded proteins and suggests that their PD-pathogenic mutations impair E3 ligase activity of the complex, which may constitute a mechanism underlying PD pathogenesis.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/parkin-pink1-and-dj-1-form-ubiquitin-e3-ligase-complex-promoting-unfolded>